

LA-UR-17-27949

Approved for public release; distribution is unlimited.

Title: Synthesis of Actinide Materials for the Study of Basic Actinide Science and Rapid Separation of Fission Products

Author(s): Dorhout, Jacquelyn Marie

Intended for: Thesis defense presentation

Issued: 2017-09-05

Disclaimer:

Los Alamos National Laboratory, an affirmative action/equal opportunity employer, is operated by the Los Alamos National Security, LLC for the National Nuclear Security Administration of the U.S. Department of Energy under contract DE-AC52-06NA25396. By approving this article, the publisher recognizes that the U.S. Government retains nonexclusive, royalty-free license to publish or reproduce the published form of this contribution, or to allow others to do so, for U.S. Government purposes. Los Alamos National Laboratory requests that the publisher identify this article as work performed under the auspices of the U.S. Department of Energy. Los Alamos National Laboratory strongly supports academic freedom and a researcher's right to publish; as an institution, however, the Laboratory does not endorse the viewpoint of a publication or guarantee its technical correctness.

Synthesis of Actinide Materials for the Study of Basic Actinide Science and Rapid Separation of Fission Products

Jacquelyn M. Dorhout

Dissertation Defense

September 29, 2017

UNCLASSIFIED

Outline

- Nuclear Forensics Story pt.1
 - Background
 - UO_2 target manufacturing
 - Irradiation and Results
- Nuclear Forensics Story pt. 2
 - MOF background
 - MOF target manufacturing
 - Irradiation and Results
- Conclusions
- Organoactinide Chemistry Story
 - Actinide Chalcogenides
 - C-X bond activation using actinides
- Summation of Projects
- Future Work
- Acknowledgements

UNCLASSIFIED

Nuclear Forensics Part 1

UO₂ Target Materials


UNCLASSIFIED

Background – Nuclear Forensics

- Nuclear forensics – evaluation of isotopic signatures to determine the identification of a device post-detonation
 - These signatures and their ratios are different depending on the type of device
 - Leads to information on when/how/where it was made
- These isotopic signatures are produced from the fission of the starting material


UNCLASSIFIED

Background - Fission



UNCLASSIFIED

Background - Fission



Background - Goal

- Goal: To create actinide target materials for the rapid separation of fission products without the need to dissolve the entire target
 - Proof-of-principle to advance the library of fission product ratios for various actinides
- Procedure: To prepare and irradiate a target material, then rapidly separate and measure the fission products

UNCLASSIFIED



Background – UO_2

- UO_2 has been used in fuel, target materials, and fission product analysis for decades
 - Literature references site these particle sizes between 20 μm and 200 μm
- Dissolution of materials is generally done in HNO_3 and/or HCl at acid concentrations $> 1\text{M}$

UNCLASSIFIED


Methods – UO₂ Production

- Hydrothermal synthesis



UNCLASSIFIED

Methods – UO₂ Production



UNCLASSIFIED


Methods – UO_2 Dissolution

- Remember – we don't want to dissolve the material
- Microparticles of UO_2 were contacted with acids to determine which would be used for future fission product extraction
 - 1M, 0.01 M HCl and 0.01 M HNO_3 tested
 - ICP-MS used to calculate concentration of uranium in acid after 24 hours
 - Unsurprisingly, lower concentrations of acids dissolved less uranium
- Acids chosen for experiments: 0.01 M, 0.1 M HCl; 0.01 M, 0.1 M HNO_3

UNCLASSIFIED

Methods – UO₂ Target Production


- KBr chosen as a secondary matrix to trap fission products
 - 3:1 ratio of KBr:UO₂ gave best homogeneity by optical spectroscopy and SEM analysis



UNCLASSIFIED

Methods – DPF Irradiation

- Dense Plasma Focus (DPF) – neutron source from D-D plasma



UNCLASSIFIED

Methods – UO₂ Target Production for DPF


- Roughly 230 mg of dUO₂ mixed with 890 mg of KBr pressed into pellets and flame sealed in glass
 - 5:1 ratio was chosen to increase the amount of material within the target
- Irradiated at DPF




UNCLASSIFIED

Results - DPF

UO₂ Solid




0.01 M HCl Solution



Methods - Irradiation



- Flattop is a critical assembly device made of HEU
- Sample holders are loaded into “Glory Hole”



UNCLASSIFIED


Methods – UO₂ Target Production for Flattop

- All targets had roughly 25 mg dUO₂ unless otherwise noted
- 3KBr:UO₂ mixture pressed into a 6 mm pellet using a KBr die
- Pellet wrapped in Al foil to prevent dispersion
- Sealed in Al sample holder




UNCLASSIFIED

Method – Rapid Separation



Methods – Rapid Separation



UNCLASSIFIED

Methods – Gamma Counting

- HPGe and BEGe detectors used in experiments
- Each sample (plus a background) was counted for the same amount of time
- Down-side: only gamma emitting isotopes are identifiable



UNCLASSIFIED

Results – Flattop Irradiations

- Several irradiations done on multiple samples
- Five UO_2 samples studied
 - Four samples of $3\text{KBr}:\text{UO}_2$
 - Each contacted with a different acid
 - 0.01 M HCl, 0.1 M HCl, 0.01 M HNO_3 , or 0.1 M HNO_3
 - One sample with no KBr present
 - How important is secondary matrix?
 - Contacted with 0.01 M HNO_3




UNCLASSIFIED

Results – Flattop Irradiations



amos
LABORATORY
943

Percent Extraction of Certain Fission Products



- Each target able to allow for extraction of a wide variety of fission products

- 0.1 M HNO₃ data is artificially high

Nuclide

- Black bars represent no KBr – secondary matrix does have an affect

UNCLASSIFIED

Issues – Part 1

- KBr is easy to work with and remove with dilute acid, but it is activated in the neutron flux to give ^{82}Br
 - Many gamma energies
 - Short half-life, high activity
 - Could be resolved by using a different secondary matrix
- Targets were irradiated at different times under different conditions
 - Cannot compare activities extracted
 - Must compare percent extraction

UNCLASSIFIED

Conclusions – Part 1

- Proof-of-concept shown
- 0.01 M HNO_3 chosen as extractant for all future experiments
- Secondary matrix does seem to have a positive affect on extraction
- Different secondary matrix could make analysis easier

UNCLASSIFIED



Nuclear Forensics Part 2

Alternative Target Materials

UNCLASSIFIED

Background – Metal Organic Frameworks

- MOFs are a porous material where metal centers are linked with organic ligands to form 3D structures
- Most common ligands are dicarboxylic acid based
- Can pores be advantageous for extracting fission products without a secondary matrix?







- Four MOFs were studied

UNCLASSIFIED



Background – UO_2 -2,6-pyridinedicarboxylic acid

- UO_2 -2,6-pydc




Immirzi, A.; Bombieri, G.; Degetto, S.; Marangoni, G. *Acta Crystallographica Section B* **1975**, 31, 1023.

Background – UO_2 -2,5-pydc





Grey = Uranium
Black = Carbon
Blue = Nitrogen
Red = Oxygen



UNCLASSIFIED




Background – UO_2 -2,4-pydc



UNCLASSIFIED

Background – UO_2 -pyromellitic acid

- Fourth MOF (UO_2 -prma) was a different structure



Mihalcea, I.; Henry, N.; Volkringer, C.; Loiseau, T. *Crystal Growth & Design* **2012**, 12, 526.

UNCLASSIFIED

Irradiation


- All four MOFs irradiated using Flattop
- Each target was contacted with 0.01 M HNO_3 in the same method as the UO_2 targets
- Prior to irradiation, a sample of each MOF was contacted in acid for 24 hours
 - Any dissolution would be due to irradiation, not material property

UNCLASSIFIED

Results - MOFs



Comparison of Extraction of Fission Products from MOFs





- Smaller pore size leads to less extraction
- UO_2 -prma target had different structure and shows best extraction for the most variety of fission products

UNCLASSIFIED

Background - Oxalate

- UO_2 -oxalate is often formed in the processes used to manufacture fuel
- Not technically a MOF, the material is a series of flat polymer sheets linked together with hydrogen bonding



UNCLASSIFIED

Results - Oxalate


- Yellow UO_2 -oxalate target returned as black UO_2
- No secondary matrix
- Some assumptions:
 - UO_2 is homogenous
 - UO_2 particles are larger than $10\text{ }\mu\text{m}$

UNCLASSIFIED

Results - Oxalate



Comparison of Fission Product Extraction from Microparticle (<10 μm)
 UO_2 and Macroparticle (>10 μm) UO_2



- Compared to microparticle UO_2 target without KBr
- Both contacted with 0.01 M HNO_3
- Microparticles are better for extraction of fission products

UNCLASSIFIED

Issues – Part 2

- Conditions in Flattop (i.e. temperature) are not ideal for using organic-based materials
- Slight enrichment of materials allows for the use of thermal neutrons (~ 0.025 eV)
 - May negate temperature issues

UNCLASSIFIED

Conclusions – Part 2

- MOFs can be used for the extraction of fission products, but yield depends on framework type and pore size
- Studying more types of MOFs will give a better correlation
- Some materials are not suited for fast neutron environments and would be better studied in a reactor situation.

UNCLASSIFIED

Organoactinide Chemistry

Actinide Chalcogenides

UNCLASSIFIED


Background

- Chalcogenides (i.e. sulfur, selenium, tellurium) are soft-donor atoms
 - Can be used for separation between lathanides and actinides
- Chemistry of the actinides (hard-donor atoms) with the soft chalcogenides is not well understood but is of interest to both the radiochemical and organoactinide communities.
 - How do 5f-electrons get involved in bonding?
 - Comparisons of thorium (no f-electrons) with uranium can help elucidate

UNCLASSIFIED

Background


- Sattelberger et.al. published the first organothorium-sulfide in 1986





UNCLASSIFIED

Experimental

- Kiplinger group was studying insertion of chalcogenides into An-C bonds and found that $(C_5Me_5)_2ThS_5$ is a thermodynamic sink




Experimental




UNCLASSIFIED

Experimental



Results

- Monitoring the reaction of $(\text{C}_5\text{Me}_5)_2\text{ThMe}_2 + \text{S}_8$ by NMR showed intermediate formation of $(\text{C}_5\text{Me}_5)_2\text{Th}(\text{SMe})_2$




Results

- Formation of MeSSMe, MeSMe, and MeSSSSMe is also seen by NMR
- $(C_5Me_5)_2Th(SMe)_2$ appears to undergo side reactions with these byproducts, but final product is always $(C_5Me_5)_2ThS_5$

UNCLASSIFIED

Results



UNCLASSIFIED

Conclusions

- There is sulfur insertion between thorium and carbon of methyl group
- Novel complex $(C_5Me_5)_2Th(SMe)_2$ reported and characterized
- Uranium chemistry doesn't behave analogously

UNCLASSIFIED

Organoactinide Chemistry

C – X Activation


UNCLASSIFIED

Background


- Transition metal mediated C–X activation ($X = \text{H}, \text{F}, \text{Cl}, \text{Br}, \text{I}$) has been studied as a route to novel organometallic compounds
- Actinide studies are lacking in the literature
 - There are examples of C–H activation with uranium
 - C–N bond activation with thorium-bipyridine complexes
- Important to understand how actinides can interact with halide environments

UNCLASSIFIED

Experimental




Experimental







UNCLASSIFIED

Results




UNCLASSIFIED

Results





Experimental




UNCLASSIFIED

Results




UNCLASSIFIED

Experimental




UNCLASSIFIED

Results



Experimental - Thorium



- Is there f-electron involvement that stabilizes the reaction?
- Theoretical calculations are in progress

UNCLASSIFIED

Conclusions

- Series of U-X (X = F, Cl, Br, I) complexes were formed using actinide mediated C–X activation
 - The benzyl group adds across the bipyridine rings
- Preliminary exploration into the mechanism seems to show that it is a migration rather than addition of the benzyl
- Chemistry does not work for thorium analog

UNCLASSIFIED

Summary and Future Work

UNCLASSIFIED

Summary – Part 1

- Proof-of-concept that an actinide target can be used to rapidly separate fission products without dissolving the target
 - Use of a secondary matrix (i.e. KBr) increases the yield

Future Work – Part 1

- Use a secondary matrix that is not activated
- Use a different isotope, or combinations of isotopes
- Use of $^{237}\text{NpO}_2$ to form ^{238}Pu

UNCLASSIFIED

Summary – Part 2

- MOFs can also be a useful target, depending on pore size and framework structure
- Use of smaller particles of UO_2 increases yield

Future Work – Part 2

- Study a variety of other MOFs
- Change the metal
- Enrich sample and use thermal neutrons

UNCLASSIFIED

Summary – Organoactinide Story

- Novel thorium sulfide
- Mechanism to understand $(C_5Me_5)_2ThS_5$
- C–X activation using uranium to give a suite of novel uranium-halide compounds
- Rare example of a U–F bond
- Rare example of activation and addition across a bipyridine ligand

UNCLASSIFIED

Future Work – Organoactinide Story

- Synthesis of $(C_5Me_5)_2US_5$ for comparison to thorium
- Theoretical explorations into C–X activation mechanism

UNCLASSIFIED

Acknowledgements



- Thesis Committee
 - Dr. Ken Czerwinski
 - Dr. Ralf Sudowe
 - Dr. Paul Forster
 - Dr. Jacqueline Kiplinger
 - Dr. Alex Barzilov
- UNLV
 - Dr. Thomas Hartmann
 - Dr. Dan Koury
 - Julie Bertoia
 - Trevor Low
 - Daniel Mast
 - Dr. Daniel Lowe, Dr. Bill Kerlin, Rebecca Springs, Lucas Boron-Brenner
 - Radiation Safety Staff
 - Radiochemistry group
- LANL
 - Drs. Karla Erickson, Justin Pagano, Stephen Cope, Ross Beattie, Marisa Monreal, David Morris, Jacqueline Veauthier, George Stanley
 - Drs. Nick Travia, Kevin Browne, Alex Lichtscheidl
- DAF
 - Dr. Todd Bredeweg
 - Dr. Donnette Lewis
 - Radiation and Safety Staff
- Family and friends
- Funding
 - NTNFC fellowship
 - Glenn T. Seaborg Institute for Transactinium Science GRA Fellowship
 - DOE Office of Basic Science



This material is based upon work supported by the U.S. Department of Homeland Security under Grant Award Number, 2012-DN-130-NF0001-02.

The views and conclusions contained in this document are those of the authors and should not be interpreted as necessarily representing the official policies, either expressed or implied, of the U.S. Department of Homeland Security.

References

- Prelas, M. A.; Weaver, C. L.; Watermann, M. L.; Lukosi, E. D.; Schott, R. J.; Wisniewski, D. A. *Progress in Nuclear Energy* **2014**, 75, 117
- Blakemore, P.; Oregon State University: Corvallis, Oregon, 2017; Vol. 2017, p image of parr bomb.
- Krishnan, M. *IEEE Transactions on Plasma Science* **2012**, 40, 3189
- Brewer, R. W.; McLaughlin, T. P.; Dean, V. *Uranium-235 Sphere Reflected by Normal Uranium Using Flattop*, Nuclear Energy Agency, 1999.
- Kim, J.; Chen, B.; Reineke, T. M.; Li, H.; Eddaoudi, M.; Moler, D. B.; O'Keeffe, M.; Yaghi, O. M. *Journal of the American Chemical Society* **2001**, 123, 8239.
- Immirzi, A.; Bombieri, G.; Degetto, S.; Marangoni, G. *Acta Crystallographica Section B* **1975**, 31, 1023.
- Mihalcea, I.; Henry, N.; Volkringer, C.; Loiseau, T. *Crystal Growth & Design* **2012**, 12, 526.
- Giesting Paul, A.; Porter Nathan, J.; Burns Peter, C. In *Zeitschrift für Kristallographie - Crystalline Materials* 2006; Vol. 221, p 252.
- Bennett Brian, K.; Harrison, R. G.; Richmond, T. G. *Journal of the American Chemical Society* **1994**, 116, 11165.
- Browne, K. P.; Maerzke, K. A.; Travia, N. E.; Morris, D. E.; Scott, B. L.; Henson, N. J.; Yang, P.; Kiplinger, J. L.; Veauthier, J. M. *Inorganic Chemistry* **2016**, 55, 4941.

UNCLASSIFIED